

[Sign In / Sign Up \(/user/login\)](/user/login)[Submit \(https://susy.mdpi.com/user/manuscripts/upload?journal=cimb\)](https://susy.mdpi.com/user/manuscripts/upload?journal=cimb)

Search for Articles:

Advanced Search

[Journals \(/about/journals\)](/about/journals) / [CIMB \(/journal/cimb\)](/journal/cimb) / [Volume 44 \(/1467-3045/44\)](/1467-3045/44) / [Issue 3 \(/1467-3045/44/3\)](/1467-3045/44/3) / [10.3390/cimb44030073](https://doi.org/10.3390/cimb44030073) /



*current issues in
molecular biology*

[\(/journal/cimb\)](/journal/cimb)

Submit to this Journal
([https://susy.mdpi.com/user/manuscripts/upload?
form%5Bjournal_id%5D%3D497](https://susy.mdpi.com/user/manuscripts/upload?form%5Bjournal_id%5D%3D497))

Review for this Journal
(<https://susy.mdpi.com/volunteer/journals/review>)

Edit a Special Issue
(</journalproposals/sendproposalspecialissue/cimb>)

We use cookies on our website to ensure you get the best experience.

Read more about our cookies [here \(/about/privacy\)](/about/privacy).

► [Article Menu](#)

Article Menu

[Accept \(/accept_cookies\)](/accept_cookies)

- [Introduction](#)
- [Materials and Methods](#)
- [Results](#)
- [Discussion](#)
- [Conclusions](#)
- [Author Contributions](#)
- [Funding](#)
- [Institutional Review Board Statement](#)
- [Informed Consent Statement](#)
- [Data Availability Statement](#)
- [Acknowledgments](#)
- [Conflicts of Interest](#)
- [References](#)

I<

Open Access Article


Intracellular Reverse Transcription of Pfizer BioNTech COVID-19 mRNA Vaccine BNT162b2 In Vitro in Human Liver Cell Line

by

 Markus Aldén (<https://sciprofiles.com/profile/1930500>) ¹  (mailto:please_login) 
(<https://orcid.org/0000-0002-5166-0055>),

 Francisco Olofsson Palla (<https://sciprofiles.com/profile/author/VTV3VHIJYWl5anhoc>)
¹  (mailto:please_login), [Read more about our cookies here \(/about/privacy\)](#).

 Daowei Yang (<https://sciprofiles.com/profile/author/LzR2bkIMamFJWtiwU0JBZkFYOI>)
¹  (mailto:please_login), [Accept \(/accept_cookies\)](#)



0

 **Mohammad Barghouth** (<https://sciprofiles.com/profile/author/c1B1b29EMVIPMWVtd3f>)
1  (mailto:please_login),
 **Cheng Luan** (<https://sciprofiles.com/profile/2059919>)¹  (mailto:please_login),
 **Magnus Rasmussen** (<https://sciprofiles.com/profile/author/a2xDSFIVNIRvaWxMVGSy>)
2  (mailto:please_login) and
 **Yang De Marinis** (<https://sciprofiles.com/profile/1713015>)^{1,*}  (mailto:please_login)
 (<https://orcid.org/0000-0002-8081-2142>)

¹ Department of Clinical Sciences, Lund University, 20502 Malmö, Sweden

² Infection Medicine, Department of Clinical Sciences, Lund University, 22362 Lund, Sweden

* Author to whom correspondence should be addressed.

Academic Editor: Stephen Malnick

Curr. Issues Mol. Biol. **2022**, *44*(3), 1115–1126; <https://doi.org/10.3390/cimb44030073>
(registering DOI)

**Received: 18 January 2022 / Revised: 19 February 2022 / Accepted: 23 February 2022 /
Published: 25 February 2022**

(This article belongs to the Topic **Clinical, Translational and Basic Research on Liver
Diseases (/topics/CTLTD)**)

[Download PDF \(/1467-3045/44/3/73/pdf\)](#)

[Browse Figures](#)

[\(/cimb/cimb-44-](#)

[00073/article_deploy/html/images/cimb-44-00073-g001.png\) \(/cimb/cimb-44-](#)

[00073/article_deploy/html/images/cimb-44-00073-g002.png\) \(/cimb/cimb-44-](#)

[00073/article_deploy/html/images/cimb-44-00073-g003.png\) \(/cimb/cimb-44-](#)

[00073/article_deploy/html/images/cimb-44-00073-g004.png\) \(/cimb/cimb-44-](#)

[00073/article_deploy/html/images/cimb-44-00073-g005.png\)](#)

[Citation Export](#)

Abstract

We use cookies on our website to ensure you get the best experience.
Read more about our cookies [here \(/about/privacy\)](#).

[Accept \(/accept_cookies\)](#)

MDPI (1)

Preclinical studies of COVID-19 mRNA vaccine BNT162b2, developed by Pfizer and BioNTech, showed reversible hepatic effects in animals that received the BNT162b2 injection. Furthermore, a recent study showed that SARS-CoV-2 RNA can be reverse-transcribed and integrated into the genome of human cells. In this study, we investigated the effect of BNT162b2 on the human liver cell line Huh7 in vitro. Huh7 cells were exposed to BNT162b2, and quantitative PCR was performed on RNA extracted from the cells. We detected high levels of BNT162b2 in Huh7 cells and changes in gene expression of long interspersed nuclear element-1 (LINE-1), which is an endogenous reverse transcriptase. Immunohistochemistry using antibody binding to LINE-1 open reading frame-1 RNA-binding protein (ORFp1) on Huh7 cells treated with BNT162b2 indicated increased nucleus distribution of LINE-1. PCR on genomic DNA of Huh7 cells exposed to BNT162b2 amplified the DNA sequence unique to BNT162b2. Our results indicate a fast up-take of BNT162b2 into human liver cell line Huh7, leading to changes in LINE-1 expression and distribution. We also show that BNT162b2 mRNA is reverse transcribed intracellularly into DNA in as fast as 6 h upon BNT162b2 exposure.

Keywords: COVID-19 mRNA vaccine (/search?q=COVID-19%20mRNA%20vaccine); BNT162b2 (/search?q=BNT162b2); liver (/search?q=liver); reverse transcription (/search?q=reverse%20transcription); LINE-1 (/search?q=LINE-1); Huh7 (/search?q=Huh7)

1. Introduction

Coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was announced by the World Health Organization (WHO) as a global pandemic on 11 March 2020, and it emerged as a devastating health crisis. As of February 2022, COVID-19 has led to over 430 million reported infection cases and 5.9 million deaths worldwide [1]. Effective and safe vaccines are urgently needed to reduce the morbidity and mortality rates associated with COVID-19.

Several vaccines for COVID-19 have been developed, with particular focus on mRNA vaccines (by Pfizer-BioNTech and Moderna), replication-defective recombinant adenoviral vector vaccines (by Janssen-Johnson and Johnson, Astra-Zeneca, Sputnik-V, and CanSino), and inactivated vaccines (by Sinopharm, Bharat Biotech and Sinovac). The mRNA vaccine has the advantages of being flexible and efficient in immunogen design and manufacturing, and currently, numerous vaccine candidates are in various stages of development and application. Specifically, COVID-19 mRNA vaccine BNT162b2 developed by Pfizer and BioNTech has been evaluated in successful clinical trials [2,3,4] and administered in national COVID-19 vaccination campaigns in different regions around the world [5,6,7,8].

BNT162b2 is a lipid nanoparticle (LNP)-encapsulated, nucleoside-modified RNA vaccine (modRNA) and encodes the full-length of SARS-CoV-2 spike (S) protein, modified by two

We use cookies on our website to ensure you get the best experience.
Read more about our cookies here (/about/privacy).

Accept (/accept_cookies)

MDPI (0)

proline mutations to ensure antigenically optimal pre-fusion conformation, which mimics the intact virus to elicit virus-neutralizing antibodies [3]. Consistent with randomized clinical trials, BNT162b2 showed high efficiency in a wide range of COVID-19-related outcomes in a real-world setting [5]. Nevertheless, many challenges remain, including monitoring for long-term safety and efficacy of the vaccine. This warrants further evaluation and investigations. The safety profile of BNT162b2 is currently only available from short-term clinical studies. Less common adverse effects of BNT162b2 have been reported, including pericarditis, arrhythmia, deep-vein thrombosis, pulmonary embolism, myocardial infarction, intracranial hemorrhage, and thrombocytopenia [4,9,10,11,12,13,14,15,16,17,18,19,20]. There are also studies that report adverse effects observed in other types of vaccines [21,22,23,24]. To better understand mechanisms underlying vaccine-related adverse effects, clinical investigations as well as cellular and molecular analyses are needed.

A recent study showed that SARS-CoV-2 RNAs can be reverse-transcribed and integrated into the genome of human cells [25]. This gives rise to the question of if this may also occur with BNT162b2, which encodes partial SARS-CoV-2 RNA. In pharmacokinetics data provided by Pfizer to European Medicines Agency (EMA), BNT162b2 biodistribution was studied in mice and rats by intra-muscular injection with radiolabeled LNP and luciferase modRNA. Radioactivity was detected in most tissues from the first time point (0.25 h), and results showed that the injection site and the liver were the major sites of distribution, with maximum concentrations observed at 8–48 h post-dose [26]. Furthermore, in animals that received the BNT162b2 injection, reversible hepatic effects were observed, including enlarged liver, vacuolation, increased gamma glutamyl transferase (γ GT) levels, and increased levels of aspartate transaminase (AST) and alkaline phosphatase (ALP) [26]. Transient hepatic effects induced by LNP delivery systems have been reported previously [27,28,29,30], nevertheless, it has also been shown that the empty LNP without modRNA alone does not introduce any significant liver injury [27]. Therefore, in this study, we aim to examine the effect of BNT162b2 on a human liver cell line in vitro and investigate if BNT162b2 can be reverse transcribed into DNA through endogenous mechanisms.

2. Materials and Methods

2.1. Cell Culture

Huh7 cells (JCRB Cell Bank, Osaka, Japan) were cultured in 37 °C at 5% CO₂ with DMEM medium (HyClone, HYCLSH30243.01) supplemented with 10% (v/v) fetal bovine serum (Sigma-Aldrich, F7524-500ML, Burlington, MA, USA) and 1% (v/v) Penicillin-Streptomycin (HyClone, SV30010, Logan, UT, USA). For BNT162b2 treatment, Huh7 cells were seeded with a density of 200,000 cells/well in 24-well plates. BNT162b2 mRNA vaccine (Pfizer BioNTech, New York, NY, USA) was diluted with sterile 0.9% sodium chloride injection, USP into a final concentration of 100 µg/mL as described in the manufacturer's guideline [31].

We use cookies on our website to ensure you get the best experience.
Read more about our cookies [here \(about/privacy\)](#).

Accept (/accept_cookies)

BNT162b2 suspension was then added in cell culture media to reach final concentrations of 0.5, 1.0, or 2.0 $\mu\text{g/mL}$. Huh7 cells were incubated with or without BNT162b2 for 6, 24, and 48 h. Cells were washed thoroughly with PBS and harvested by trypsinization and stored in -80°C until further use.

2.2. REAL-TIME RT-QPCR

RNA from the cells was extracted with RNeasy Plus Mini Kit (Qiagen, 74134, Hilden, Germany) following the manufacturer's protocol. RT-PCR was performed using RevertAid First Strand cDNA Synthesis kit (Thermo Fisher Scientific, K1622, Waltham, MA, USA) following the manufacturers protocol. Real-time qPCR was performed using Maxima SYBR Green/ROX qPCR Master Mix (Thermo Fisher Scientific, K0222, Waltham, MA, USA) with primers for BNT162b2, *LINE-1* and housekeeping genes *ACTB* and *GAPDH* (Table 1).

Table 1. Primer sequences of RT-qPCR and PCR.

2.3. Immunofluorescence Staining and Confocal Imaging

Huh7 cells were cultured in eight-chamber slides (LAB-TEK, 154534, Santa Cruz, CA, USA) with a density of 40,000 cells/well, with or without BNT162b2 (0.5, 1 or 2 $\mu\text{g/mL}$) for 6 h. Immunohistochemistry was performed using primary antibody anti-LINE-1 ORF1p mouse monoclonal antibody (Merck, 3574308, Kenilworth, NJ, USA), secondary antibody Cy3 Donkey anti-mouse (Jackson ImmunoResearch, West Grove, PA, USA), and Hoechst (Life technologies, 34850, Carlsbad, CA, USA), following the protocol from Thermo Fisher (Waltham, MA, USA). Two images per condition were taken using a Zeiss LSM 800 and a 63X oil immersion objective, and the staining intensity was quantified on the individual whole cell area and the nucleus area on 15 cells per image by ImageJ 1.53c. LINE-1 staining intensity for the cytosol was calculated by subtracting the intensity of the nucleus from that of the whole cell. All images of the cells were assigned a random number to prevent bias. To mark the nuclei (determined by the Hoechst staining) and the whole cells (determined by the borders of the LINE-1 fluorescence), the Freehand selection tool was used. These areas were then measured, and the mean intensity was used to compare the groups.

We use cookies on our website to ensure you get the best experience.

2.4. Genomic DNA Purification, PCR Amplification, Agarose Gel Purification, and Sanger Sequencing

Genomic DNA was extracted from cell pellets with PBD buffer (10 mM Tris-HCl pH 8.3, 50 mM KCl, 2.5 mM MgCl_2 , 0.45% NP-40, 0.45% Tween-20) according to protocol described

previously [32]. To remove residual RNA from the DNA preparation, RNase (100 µg/mL, Qiagen, Hilden, Germany) was added to the DNA preparation and incubated at 37 °C for 3 h, followed by 5 min at 95 °C. PCR was then performed using primers targeting BNT162b2 (sequences are shown in **Table 1**), with the following program: 5 min at 95 °C, 35 cycles of 95 °C for 30 s, 58 °C for 30 s, and 72 °C for 1 min; finally, 72 °C for 5 min and 12 °C for 5 min. PCR products were run on 1.4% (w/v) agarose gel. Bands corresponding to the amplicons of the expected size (444 bps) were cut out and DNA was extracted using QIAquick PCR Purification Kit (Qiagen, 28104, Hilden, Germany), following the manufacturer's instructions. The sequence of the DNA amplicon was verified by Sanger sequencing (Eurofins Genomics, Ebersberg, Germany).

Statistics

Statistical comparisons were performed using two-tailed Student's *t*-test and ANOVA. Data are expressed as the mean ± SEM or ± SD. Differences with *p* < 0.05 are considered significant.

2.5. Ethical Statements

The Huh7 cell line was obtained from Japanese Collection of Research Bioresources (JCRB) Cell Bank.

3. Results

3.1. BNT162b2 Enters Human Liver Cell Line Huh7 Cells at High Efficiency

To determine if BNT162b2 enters human liver cells, we exposed human liver cell line Huh7 to BNT162b2. In a previous study on the uptake kinetics of LNP delivery in Huh7 cells, the maximum biological efficacy of LNP was observed between 4–7 h [33]. Therefore, in our study, Huh7 cells were cultured with or without increasing concentrations of BNT162b2 (0.5, 1.0 and 2.0 µg/mL) for 6, 24, and 48 h. RNA was extracted from cells and a real-time quantitative reverse transcription polymerase chain reaction (RT-qPCR) was performed using primers targeting the BNT162b2 sequence, as illustrated in **Figure 1**. The full sequence of BNT162b2 is publicly available [34] and contains a two-nucleotides cap; 5'- untranslated region (UTR) that incorporates the 5' -UTR of a human α-globin gene; the full-length of SARS-CoV-2 S protein with two proline mutations; 3'-UTR that incorporates the human mitochondrial 12S rRNA (mtRNR1) segment and human AES/TLE5 gene segment with two C→U mutations; poly(A) tail. Detailed analysis of the S protein sequence in BNT162b2

We use cookies on our website to ensure you get the best experience.
Read more about our cookies [here \(about/privacy\)](#).
revealed 124 sequences that are 100% identical to human genomic sequences and three sequences with only one nucleotide (nt) mismatch in 19–26 nts (**Table S1, see Supplementary Materials**). To detect BNT162b2 RNA level, we designed primers with forward primer located in SARS-CoV-2 S protein regions and reverse primer in 3'-UTR, which

allows detection of PCR amplicon unique to BNT162b2 without unspecific binding of the primers to human genomic regions.




Figure 1. PCR primer set used to detect mRNA level and reverse-transcription of BNT162b2. Illustration of BNT162b2 was adapted from previously described literature [34].

RT-qPCR results showed that Huh7 cells treated with BNT162b2 had high levels of BNT162b2 mRNA relative to housekeeping genes at 6, 24, and 48 h (**Figure 2**, presented in logged $2^{-\Delta\Delta CT}$ due to exceptionally high levels). The three BNT162b2 concentrations led to similar intracellular BNT162b2 mRNA levels at the different time points, except that the significant difference between 1.0 and 2.0 $\mu\text{g/mL}$ was observed at 48 h. BNT162b2 mRNA levels were significantly decreased at 24 h compared to 6 h, but increased again at 48 h.




Figure 2. BNT162b2 mRNA levels in Huh7 cells treated with BNT162b2. Huh7 cells were treated without (Ctrl) or with 0.5 (V1), 1 (V2), and 2 $\mu\text{g/mL}$ (V3) of BNT162b2 for 6 (green dots), 24 (orange dots), and 48 h (blue dots). RNA was purified and qPCR was performed using primers targeting BNT162b2. RNA levels of BNT162b2 are presented as logged $2^{-\Delta\Delta CT}$ values relative to house-keeping genes *GAPDH* and *ACTB*. Results are from five independent experiments ($n = 5$). Differences between respective groups were analyzed using two-tailed Student's *t*-test. Data are expressed as the mean $\pm$ SEM. (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ vs. respective control at each time point, or as indicated).

3.2 Effect of BNT162b2 on Human Endogenous Reverse Transcriptase Long Interspersed Nuclear Element-1 (*LINE-1*)

Here we examined the effect of BNT162b2 on *LINE-1* gene expression. RT-qPCR was performed on RNA purified from Huh7 cells treated with BNT162b2 (0, 0.5, 1.0, and 2.0 $\mu\text{g/mL}$) for 6, 24, and 48 h, using primers targeting *LINE-1*. Significantly increased *LINE-1* expression compared to control was observed at 6 h by 2.0 $\mu\text{g/mL}$ BNT162b2, while lower BNT162b2 concentrations decreased *LINE-1* expression at all time points (**Figure 3**).




Figure 3. *LINE-1* mRNA levels in Huh7 cells treated with BNT162b2. Huh7 cells were treated without (Ctrl) or with 0.5 (V1), 1 (V2), and 2 $\mu\text{g/mL}$ (V3) of BNT162b2 for 6 (green dots), 24 (red dots), and 48 h (blue dots). RNA was purified and qPCR was performed using primers targeting *LINE-1*. RNA levels of *LINE-1* are presented as $2^{-\Delta\Delta\text{CT}}$ values relative to house-keeping genes *GAPDH* and *ACTB*. Results are from five independent experiments ($n = 5$). Differences between respective groups were analyzed using two-tailed Student's *t*-test. Data are expressed as the mean $\pm$ SEM. (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ vs. respective control at each time point, or as indicated; † $p < 0.05$ vs. 6 h-Ctrl).

Next, we studied the effect of BNT162b2 on *LINE-1* protein level. The full-length *LINE-1* consists of a 5' untranslated region (UTR), two open reading frames (ORFs), ORF1 and ORF2, and a 3'UTR, of which ORF1 is an RNA binding protein with chaperone activity. The retrotransposition activity of *LINE-1* has been demonstrated to involve ORF1 translocation to the nucleus [35]. Huh7 cells treated with or without BNT162b2 (0.5, 1.0 and 2.0 $\mu\text{g/mL}$) for 6 h were fixed and stained with antibodies binding to *LINE-1* ORF1p, and DNA-specific probe Hoechst for visualization of cell nucleus (**Figure 4a**). Quantification of immunofluorescence staining intensity showed that BNT162b2 increased *LINE-1* ORF1p protein levels in both the whole cell area and nucleus at all concentrations tested (**Figure 4b–d**).




Figure 4. Immunohistochemistry of Huh7 cells treated with BNT162b2 on LINE-1 protein distribution. Huh7 cells were treated without (Ctrl) or with 0.5, 1, and 2 µg/mL of BNT162b2 for 6 h. Cells were fixed and stained with antibodies binding to LINE-1 ORF1p (red) and DNA-specific probe Hoechst for visualization of cell nucleus (blue). (a) Representative images of LINE-1 expression in Huh7 cells treated with or without BNT162b2. (b–d) Quantification of LINE-1 protein in whole cell area (b), cytosol (c), and nucleus (d). All data were analyzed using One-Way ANOVA, and graphs were created using GraphPad Prism V 9.2. All data is presented as mean ± SD (** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$ as indicated).

3.3. Detection of Reverse Transcribed BNT162b2 DNA in Huh7 Cells

A previous study has shown that entry of LINE-1 protein into the nucleus is associated with retrotransposition [35]. In the immunofluorescence staining experiment described above, increased levels of LINE-1 in the nucleus were observed already at the lowest concentration of BNT162b2 (0.5 µg/mL). To examine if BNT162b2 is reversely transcribed into DNA when LINE-1 is elevated, we purified genomic DNA from Huh7 cells treated with 0.5 µg/mL of BNT162b2 for 6, 24, and 48 h. Purified DNA was treated with RNase to remove RNA and subjected to PCR using primers targeting BNT162b2, as illustrated in **Figure 1**. Amplified DNA fragments were then visualized by electrophoresis and gel-purified (**Figure 5**). BNT162b2 DNA amplicons were detected in all three time points (6, 24, and 48 h). Sanger

We use cookies on our website to ensure you get the best experience.
Read more about our cookies here (about/privacy).
sequencing confirmed that the DNA amplicons were identical to the BNT162b2 sequence flanked by the primers (**Table 2**). To ensure that the DNA amplicons were derived from DNA but not BNT162b2 RNA, we also performed PCR on RNA purified from Huh7 cells treated

with 0.5 $\mu\text{g/mL}$ BNT162b2 for 6 h, with or without RNase treatment (Ctrl 5 and 6 in **Figure 5**), and no amplicon was detected in the RNA samples subjected to PCR.




Figure 5. Detection of DNA amplicons of BNT162b2 in Huh7 cells treated with BNT162b2. Huh7 cells were treated without (Ctrl) or with 0.5 $\mu\text{g/mL}$ of BNT162b2 for 6, 24, and 48 h. Genomic DNA was purified and digested with 100 $\mu\text{g/mL}$ RNase. PCR was run on all samples with primers targeting BNT162b2, as shown in **Figure 1** and **Table 1**. DNA amplicons (444 bps) were visualized on agarose gel. BNT: BNT162b2; L: DNA ladder; Ctrl1: cultured Huh7 cells; Ctrl2: Huh7 cells without BNT162b2 treatment collected at 6 h; Ctrl3: Huh7 cells without BNT162b2 treatment collected at 24 h; Ctrl4: Huh7 cells without BNT162b2 treatment collected at 48 h; Ctrl5: RNA from Huh7 cells treated with 0.5 $\mu\text{g/mL}$ of BNT162b2 for 6 h; Ctrl6: RNA from Huh7 cells treated with 0.5 $\mu\text{g/mL}$ of BNT162b2 for 6 h, digested with RNase.

Table 2. Sanger sequencing result of the BNT162b2 amplicon.

4. Discussion

In this study we present evidence that COVID-19 mRNA vaccine BNT162b2 is able to enter the human liver cell line Huh7 in vitro. BNT162b2 mRNA is reverse transcribed intracellularly into DNA as fast as 6 h after BNT162b2 exposure. A possible mechanism for reverse transcription is through endogenous reverse transcriptase LINE-1, and the nucleus protein distribution of LINE-1 is elevated by BNT162b2.

We use cookies on our website to ensure you get the best experience. Read more about our cookies [here](#) (about privacy). Intracellular accumulation of LNP in hepatocytes has been demonstrated in vivo [36]. A preclinical study on BNT162b2 showed that BNT162b2 enters the human cell line HEK293T cells and leads to robust expression of BNT162b2 antigen [37]. Therefore, in this study, we first investigated the entry of BNT162b2 in the human liver cell line Huh7 cells. The choice of

MDPI (J)

BNT162b2 concentrations used in this study warrants explanation. BNT162b2 is administered as a series of two doses three weeks apart, and each dose contains 30 µg of BNT162b2 in a volume of 0.3 mL, which makes the local concentration at the injection site at the highest 100 µg/mL [31]. A previous study on mRNA vaccines against H10N8 and H7N9 influenza viruses using a similar LNP delivery system showed that the mRNA vaccine can distribute rather nonspecifically to several organs such as liver, spleen, heart, kidney, lung, and brain, and the concentration in the liver is roughly 100 times lower than that of the intra-muscular injection site [38]. In the assessment report on BNT162b2 provided to EMA by Pfizer, the pharmacokinetic distribution studies in rats demonstrated that a relatively large proportion (up to 18%) of the total dose distributes to the liver [26]. We therefore chose to use 0.5, 1, and 2 µg/mL of vaccine in our experiments on the liver cells. However, the effect of a broader range of lower and higher concentrations of BNT162b2 should also be verified in future studies. ▶

In the current study, we employed a human liver cell line for in vitro investigation. It is worth investigating if the liver cells also present the vaccine-derived SARS-CoV-2 spike protein, which could potentially make the liver cells targets for previously primed spike protein reactive cytotoxic T cells. There has been case reports on individuals who developed autoimmune hepatitis [39] after BNT162b2 vaccination. To obtain better understanding of the potential effects of BNT162b2 on liver function, in vivo models are desired for future studies.

In the BNT162b2 toxicity report, no genotoxicity nor carcinogenicity studies have been provided [26]. Our study shows that BNT162b2 can be reverse transcribed to DNA in liver cell line Huh7, and this may give rise to the concern if BNT162b2-derived DNA may be integrated into the host genome and affect the integrity of genomic DNA, which may potentially mediate genotoxic side effects. At this stage, we do not know if DNA reverse transcribed from BNT162b2 is integrated into the cell genome. Further studies are needed to demonstrate the effect of BNT162b2 on genomic integrity, including whole genome sequencing of cells exposed to BNT162b2, as well as tissues from human subjects who received BNT162b2 vaccination.

Human autonomous retrotransposon LINE-1 is a cellular endogenous reverse transcriptase and the only remaining active transposon in humans, able to retrotranspose itself and other nonautonomous elements [40,41], and ~17% of the human genome are comprised of LINE-1 sequences [42]. The nonautonomous *Alu* elements, short, interspersed nucleotide elements (SINEs), variable-number-of-tandem-repeats (VNTR), as well as cellular mRNA-processed pseudogenes, are retrotransposed by the LINE-1 retrotransposition proteins working in *trans* [43,44]. A recent study showed that endogenous LINE-1 mediates reverse transcription and integration of SARS-CoV-2 sequences in the genomes of infected human cells [25]. Furthermore, expression of endogenous LINE-1 is often increased upon viral infection, including SARS-CoV-2 infection [45,46,47]. Previous studies showed that LINE-1 retrotransposition activity is regulated by RNA metabolism [48,49], DNA damage response [50], and autophagy [51]. Efficient retrotransposition of LINE-1 is often associated with cell cycle and nuclear envelope breakdown during mitosis [52,53], as well as exogenous

We use cookies on our website to ensure you get the best experience.
Read information about our cookies here (about privacy).

Accept (accept_cookies)

MDPI (0)

retroviruses [54,55], which promotes entrance of LINE-1 into the nucleus. In our study, we observed increased LINE-1 ORF1p distribution as determined by immunohistochemistry in the nucleus by BNT162b2 at all concentrations tested (0.5, 1, and 2 µg/mL), while elevated *LINE-1* gene expression was detected at the highest BNT162b2 concentration (2 µg/mL). It is worth noting that gene transcription is regulated by chromatin modifications, transcription factor regulation, and the rate of RNA degradation, while translational regulation of protein involves ribosome recruitment on the initiation codon, modulation of peptide elongation, termination of protein synthesis, or ribosome biogenesis. These two processes are controlled by different mechanisms, and therefore they may not always show the same change patterns in response to external challenges. The exact regulation of LINE-1 activity in response to BNT162b2 merits further study.

◀ The cell model that we used in this study is a carcinoma cell line, with active DNA replication which differs from non-dividing somatic cells. It has also been shown that Huh7 cells display significant different gene and protein expression including upregulated proteins involved in RNA metabolism [56]. However, cell proliferation is also active in several human tissues such as the bone marrow or basal layers of epithelia as well as during embryogenesis, and it is therefore necessary to examine the effect of BNT162b2 on genomic integrity under such conditions. Furthermore, effective retrotransposition of LINE-1 has also been reported in non-dividing and terminally differentiated cells, such as human neurons [57,58].

The Pfizer EMA assessment report also showed that BNT162b2 distributes in the spleen (<1.1%), adrenal glands (<0.1%), as well as low and measurable radioactivity in the ovaries and testes (<0.1%) [26]. Furthermore, no data on placental transfer of BNT162b2 is available from Pfizer EMA assessment report. Our results showed that BNT162b2 mRNA readily enters Huh7 cells at a concentration (0.5 µg/mL) corresponding to 0.5% of the local injection site concentration, induce changes in LINE-1 gene and protein expression, and within 6 h, reverse transcription of BNT162b2 can be detected. It is therefore important to investigate further the effect of BNT162b2 on other cell types and tissues both in vitro and in vivo.

5. Conclusions

Our study is the first in vitro study on the effect of COVID-19 mRNA vaccine BNT162b2 on human liver cell line. We present evidence on fast entry of BNT162b2 into the cells and subsequent intracellular reverse transcription of BNT162b2 mRNA into DNA.

We use cookies on our website to ensure you get the best experience.
Supplementary Materials
Read more about our cookies [here \(/about/privacy\)](#).

The following supporting information can be downloaded at:
<https://www.mdpi.com/article/10.3390/cimb44030073/s1> **Accept (/accept_cookies)**

Author Contributions



M.A., F.O.F., D.Y., M.B. and C.L. performed in vitro experiments. M.A. and F.O.F. performed data analysis. M.R. and Y.D.M. contributed to the implementation of the research, designed, and supervised the study. Y.D.M. wrote the paper with input from all authors. All authors have read and agreed to the published version of the manuscript.

Funding

This study was supported by the Swedish Research Council, Strategic Research Area Exodiab, Dnr 2009-1039, the Swedish Government Fund for Clinical Research (ALF) and the foundation of Skåne University Hospital.

Institutional Review Board Statement

Not applicable.

Informed Consent Statement

Not applicable.

Data Availability Statement

All data supporting the findings of this study are available within the article and supporting information.

Acknowledgments

The authors thank Sven Haidl, Maria Josephson, Enming Zhang, Jia-Yi Li, Caroline Haikal, and Pradeep Bompada for their support to this study.

Conflicts of Interest

We use cookies on our website to ensure you get the best experience.

[Read more about our cookies here \(/about/privacy\).](#)

The authors declare no conflict of interest.

[Accept \(/accept_cookies\)](#)

References

2. Mulligan, M.J.; Lyke, K.E.; Kitchin, N.; Absalon, J.; Gurtman, A.; Lockhart, S.; Neuzil, K.; Raabe, V.; Bailey, R.; Swanson, K.A.; et al. Phase I/II study of COVID-19 RNA vaccine BNT162b1 in adults. *Nature* **2020**, *586*, 589–593. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Phase+I/II+study+of+COVID-19+RNA+vaccine+BNT162b1+in+adults&author=Mulligan,+M.J.&author=Lyke,+K.E.&author=Kitchin,+N.&author=Absalon,+J.&author=Gurtman,+A.&author=Lockhart,+S.&author=Neuzil,+K.&author=Raabe,+V.&author=Bailey,+R.&author=Swanson,+K.A.&publication_year=2020&journal=Nature&volume=586&pages=589%E2%80%93593&doi=10.1038/s41586-020-2639-4&pmid=32785213)] [CrossRef] (<https://doi.org/10.1038/s41586-020-2639-4>) [PubMed] (<https://www.ncbi.nlm.nih.gov/pubmed/32785213>)]
3. Walsh, E.E.; Frenck, R.W., Jr.; Falsey, A.R.; Kitchin, N.; Absalon, J.; Gurtman, A.; Lockhart, S.; Neuzil, K.; Mulligan, M.J.; Bailey, R.; et al. Safety and Immunogenicity of Two RNA-Based COVID-19 Vaccine Candidates. *N. Engl. J. Med.* **2020**, *383*, 2439–2450. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Safety+and+Immunogenicity+of+Two+RNA-Based+COVID-19+Vaccine+Candidates&author=Walsh,+E.E.&author=Frenck,+R.W.,+Jr.&author=Falsey,+A.R.&author=Kitchin,+N.&author=Absalon,+J.&author=Gurtman,+A.&author=Lockhart,+S.&author=Neuzil,+K.&author=Mulligan,+M.J.&author=Bailey,+R.&publication_year=2020&journal=N.+Engl.+J.+Med.&volume=383&pages=2439%E2%80%932450&doi=10.1056/NEJMoa2027906&pmid=33053279)] [CrossRef] (<https://doi.org/10.1056/NEJMoa2027906>) [PubMed] (<https://www.ncbi.nlm.nih.gov/pubmed/33053279>)]
4. Polack, F.P.; Thomas, S.J.; Kitchin, N.; Absalon, J.; Gurtman, A.; Lockhart, S.; Perez, J.L.; Perez Marc, G.; Moreira, E.D.; Zerbini, C.; et al. Safety and Efficacy of the BNT162b2 mRNA COVID-19 Vaccine. *N. Engl. J. Med.* **2020**, *383*, 2603–2615. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Safety+and+Efficacy+of+the+BNT162b2+mRNA+COVID-19+Vaccine&author=Polack,+F.P.&author=Thomas,+S.J.&author=Kitchin,+N.&author=Absalon,+J.&author=Gurtman,+A.&author=Lockhart,+S.&author=Perez,+J.L.&author=Perez+Marc,+G.&author=Moreira,+E.D.&author=Zerbini,+C.&publication_year=2020&journal=N.+Engl.+J.+Med.&volume=383&pages=2603%E2%80%932615&doi=10.1056/NEJMoa2027906&pmid=33053279)] [CrossRef] (<https://doi.org/10.1056/NEJMoa2027906>) [PubMed] (<https://www.ncbi.nlm.nih.gov/pubmed/33301246>)]

We used the data from the NEJMoa2027906 and NEJMoa2027906 for the analysis.

Read more at <https://doi.org/10.1056/NEJMoa2027906>.

(<https://www.ncbi.nlm.nih.gov/pubmed/33301246>)

5. Harris, R.J.; Hall, J.A.; Zaidi, A.; Andrews, N.J.; Dunbar, J.K.; Dabrera, G. Effect of Vaccination on Household Transmission of SARS-CoV-2 in England. *N. Engl. J. Med.* **2021**, *385*, 759–760. [Google Scholar]

([https://scholar.google.com/scholar_lookup?](https://scholar.google.com/scholar_lookup?title=Effect+of+Vaccination+on+Household+Transmission+of+SARS-CoV-2+in+England&author=Harris,+R.J.&author=Hall,+J.A.&author=Zaidi,+A.&author=Andrews,+N.J.&author=Dunbar,+J.K.&author=Dabrera,+G.&publication_year=2021&journal=N.+Engl.+J.+Med.&volume=385&pages=759%E2%80%93760&doi=10.1056/NEJMc2107717)

[title=Effect+of+Vaccination+on+Household+Transmission+of+SARS-CoV-2+in+England&author=Harris,+R.J.&author=Hall,+J.A.&author=Zaidi,+A.&author=Andrews,+N.J.&author=Dunbar,+J.K.&author=Dabrera,+G.&publication_year=2021&journal=N.+Engl.+J.+Med.&volume=385&pages=759%E2%80%93760&doi=10.1056/NEJMc2107717](https://scholar.google.com/scholar_lookup?title=Effect+of+Vaccination+on+Household+Transmission+of+SARS-CoV-2+in+England&author=Harris,+R.J.&author=Hall,+J.A.&author=Zaidi,+A.&author=Andrews,+N.J.&author=Dunbar,+J.K.&author=Dabrera,+G.&publication_year=2021&journal=N.+Engl.+J.+Med.&volume=385&pages=759%E2%80%93760&doi=10.1056/NEJMc2107717)) [CrossRef (<https://doi.org/10.1056/NEJMc2107717>)]

6. Butt, A.A.; Omer, S.B.; Yan, P.; Shaikh, O.S.; Mayr, F.B. SARS-CoV-2 Vaccine Effectiveness in a High-Risk National Population in a Real-World Setting. *Ann. Intern. Med.* **2021**, *174*, 1404–1408. [Google Scholar]

([https://scholar.google.com/scholar_lookup?title=SARS-CoV-](https://scholar.google.com/scholar_lookup?title=SARS-CoV-2+Vaccine+Effectiveness+in+a+High-Risk+National+Population+in+a+Real-World+Setting&author=Butt,+A.A.&author=Omer,+S.B.&author=Yan,+P.&author=Shaikh,+O.S.&author=Mayr,+F.B.&publication_year=2021&journal=Ann.+Intern.+Med.&volume=174&pages=1404%E2%80%931408&doi=10.7326/M21-1577)

[2+Vaccine+Effectiveness+in+a+High-Risk+National+Population+in+a+Real-World+Setting&author=Butt,+A.A.&author=Omer,+S.B.&author=Yan,+P.&author=Shaikh,+O.S.&author=Mayr,+F.B.&publication_year=2021&journal=Ann.+Intern.+Med.&volume=174&pages=1404%E2%80%931408&doi=10.7326/M21-1577](https://scholar.google.com/scholar_lookup?title=SARS-CoV-2+Vaccine+Effectiveness+in+a+High-Risk+National+Population+in+a+Real-World+Setting&author=Butt,+A.A.&author=Omer,+S.B.&author=Yan,+P.&author=Shaikh,+O.S.&author=Mayr,+F.B.&publication_year=2021&journal=Ann.+Intern.+Med.&volume=174&pages=1404%E2%80%931408&doi=10.7326/M21-1577))

[CrossRef (<https://doi.org/10.7326/M21-1577>)]

7. Dagan, N.; Barda, N.; Kepten, E.; Miron, O.; Perchik, S.; Katz, M.A.; Hernan, M.A.; Lipsitch, M.; Reis, B.; Balicer, R.D. BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination Setting. *N. Engl. J. Med.* **2021**, *384*, 1412–1423. [Google Scholar]

([https://scholar.google.com/scholar_lookup?](https://scholar.google.com/scholar_lookup?title=BNT162b2+mRNA+Covid-19+Vaccine+in+a+Nationwide+Mass+Vaccination+Setting&author=Dagan,+N.&author=Barda,+N.&author=Kepten,+E.&author=Miron,+O.&author=Perchik,+S.&author=Katz,+M.A.&author=Hernan,+M.A.&author=Lipsitch,+M.&author=Reis,+B.&author=Balicer,+R.D.&publication_year=2021&journal=N.+Engl.+J.+Med.&volume=384&pages=1412%E2%80%931423&doi=10.1056/NEJMoa2101765)

[title=BNT162b2+mRNA+Covid-19+Vaccine+in+a+Nationwide+Mass+Vaccination+Setting&author=Dagan,+N.&author=Barda,+N.&author=Kepten,+E.&author=Miron,+O.&author=Perchik,+S.&author=Katz,+M.A.&author=Hernan,+M.A.&author=Lipsitch,+M.&author=Reis,+B.&author=Balicer,+R.D.&publication_year=2021&journal=N.+Engl.+J.+Med.&volume=384&pages=1412%E2%80%931423&doi=10.1056/NEJMoa2101765](https://scholar.google.com/scholar_lookup?title=BNT162b2+mRNA+Covid-19+Vaccine+in+a+Nationwide+Mass+Vaccination+Setting&author=Dagan,+N.&author=Barda,+N.&author=Kepten,+E.&author=Miron,+O.&author=Perchik,+S.&author=Katz,+M.A.&author=Hernan,+M.A.&author=Lipsitch,+M.&author=Reis,+B.&author=Balicer,+R.D.&publication_year=2021&journal=N.+Engl.+J.+Med.&volume=384&pages=1412%E2%80%931423&doi=10.1056/NEJMoa2101765)) [CrossRef (<https://doi.org/10.1056/NEJMoa2101765>)]

8. Rossman, H.; Shilo, S.; Meir, T.; Gorfine, M.; Shalit, U.; Segal, E. COVID-19 dynamics after a national immunization program in Israel. *Nat. Med.* **2021**, *27*, 1055–1061. [Google Scholar]

([https://scholar.google.com/scholar_lookup?title=COVID-](https://scholar.google.com/scholar_lookup?title=COVID-19+dynamics+after+a+national+immunization+program+in+Israel&author=Rossman,+H.&author=Shilo,+S.&author=Meir,+T.&author=Gorfine,+M.&author=Shalit,+U.&author=Segal,+E.&publication_year=2021&journal=Nat.+Med.&volume=27&pages=1055%E2%80%931061&doi=10.1038/s41591-021-01337-2)

[19+dynamics+after+a+national+immunization+program+in+Israel&author=Rossman,+H.&author=Shilo,+S.&author=Meir,+T.&author=Gorfine,+M.&author=Shalit,+U.&author=Segal,+E.&publication_year=2021&journal=Nat.+Med.&volume=27&pages=1055%E2%80%931061&doi=10.1038/s41591-021-01337-2](https://scholar.google.com/scholar_lookup?title=COVID-19+dynamics+after+a+national+immunization+program+in+Israel&author=Rossman,+H.&author=Shilo,+S.&author=Meir,+T.&author=Gorfine,+M.&author=Shalit,+U.&author=Segal,+E.&publication_year=2021&journal=Nat.+Med.&volume=27&pages=1055%E2%80%931061&doi=10.1038/s41591-021-01337-2)) [CrossRef (<https://doi.org/10.1038/s41591-021-01337-2>)]

- Fan, B.E.; Shen, J.Y.; Lim, X.R.; Tu, T.M.; Chang, C.C.R.; Khin, H.S.W.; Koh, J.S.; Rao, J.P.; Lau, S.L.; Tan, G.B.; et al. Cerebral venous thrombosis post BNT162b2 mRNA SARS-CoV-2 vaccination: A black swan event. *Am. J. Hematol.* **2021**, *96*, E357–E361. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Cerebral+venous+thrombosis+post+BNT162b2+mRNA+SARS-CoV-2+vaccination:+A+black+swan+event&author=Fan,+B.E.&author=Shen,+J.Y.&author=Lim,+X.R.&author=Tu,+T.M.&author=Chang,+C.C.R.&author=Khin,+H.S.W.&author=Koh,+J.S.&author=Rao,+J.P.&author=Lau,+S.L.&author=Tan,+G.B.&publication_year=2021&journal=Am.+J.+Hematol.&volume=96&pages=E357%E2%80%93E361&doi=10.1002/ajh.26272)] [CrossRef (<https://doi.org/10.1002/ajh.26272>)]
10. Larson, K.F.; Ammirati, E.; Adler, E.D.; Cooper, L.T., Jr.; Hong, K.N.; Saponara, G.; Couri, D.; Cereda, A.; Procopio, A.; Cavalotti, C.; et al. Myocarditis After BNT162b2 and mRNA-1273 Vaccination. *Circulation* **2021**, *144*, 506–508. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Myocarditis+After+BNT162b2+and+mRNA-1273+Vaccination&author=Larson,+K.F.&author=Ammirati,+E.&author=Adler,+E.D.&author=Cooper,+L.T.,+Jr.&author=Hong,+K.N.&author=Saponara,+G.&author=Couri,+D.&author=Cereda,+A.&author=Procopio,+A.&author=Cavalotti,+C.&publication_year=2021&journal=Circulation&volume=144&pages=506%E2%80%93508&doi=10.1161/CIRCULATIONAHA.121.055913)] [CrossRef (<https://doi.org/10.1161/CIRCULATIONAHA.121.055913>)]
11. Menni, C.; Klaser, K.; May, A.; Polidori, L.; Capdevila, J.; Louca, P.; Sudre, C.H.; Nguyen, L.H.; Drew, D.A.; Merino, J.; et al. Vaccine side-effects and SARS-CoV-2 infection after vaccination in users of the COVID Symptom Study app in the UK: A prospective observational study. *Lancet Infect. Dis.* **2021**, *21*, 939–949. [Google Scholar ([https://scholar.google.com/scholar_lookup?title=Vaccine+side-effects+and+SARS-CoV-2+infection+after+vaccination+in+users+of+the+COVID+Symptom+Study+app+in+the+UK:+A+prospective+observational+study&author=Menni,+C.&author=Klaser,+K.&author=May,+A.&author=Polidori,+L.&author=Capdevila,+J.&author=Louca,+P.&author=Sudre,+C.H.&author=Nguyen,+L.H.&author=Drew,+D.A.&author=Merino,+J.&publication_year=2021&journal=Lancet+Infect.+Dis.&volume=21&pages=939%E2%80%93949&doi=10.1016/S1473-3099\(21\)00224-3](https://scholar.google.com/scholar_lookup?title=Vaccine+side-effects+and+SARS-CoV-2+infection+after+vaccination+in+users+of+the+COVID+Symptom+Study+app+in+the+UK:+A+prospective+observational+study&author=Menni,+C.&author=Klaser,+K.&author=May,+A.&author=Polidori,+L.&author=Capdevila,+J.&author=Louca,+P.&author=Sudre,+C.H.&author=Nguyen,+L.H.&author=Drew,+D.A.&author=Merino,+J.&publication_year=2021&journal=Lancet+Infect.+Dis.&volume=21&pages=939%E2%80%93949&doi=10.1016/S1473-3099(21)00224-3))] [CrossRef ([https://doi.org/10.1016/S1473-3099\(21\)00224-3](https://doi.org/10.1016/S1473-3099(21)00224-3))]

12. Hansen, T.; Titze, U.; Kulamadayil-Heidenreich, N.S.A.; Glombitza, S.; Tebbe, J.J.; Roeken, C.; Schulz, B.; Weise, M.; Wilkens, L. First case of postmortem study in a patient vaccinated against SARS-CoV-2. *Int. J. Infect. Dis.* **2021**, *107*, 172–175. [Google Scholar (https://scholar.google.com/scholar_lookup?title=First+case+of+postmortem+study+in+a+patient+vaccinated+against+SARS-CoV-2&author=Hansen,+T.&author=Titze,+U.&author=Kulamadayil-Heidenreich,+N.S.A.&author=Glombitza,+S.&author=Tebbe,+J.J.&author=Roeken,+C.&author=Schulz,+B.&author=Weise,+M.&author=Wilkens,+L.&publication_year=2021&journal=Int.+J.+Infect.+Dis.&volume=107&pages=172%E2%80%93175&doi=10.1016/j.ijid.2021.04.053&pmid=33872783)] [CrossRef] (<https://doi.org/10.1016/j.ijid.2021.04.053>) [PubMed] (<https://www.ncbi.nlm.nih.gov/pubmed/33872783>)]
13. Kadali, R.A.K.; Janagama, R.; Peruru, S.; Malayala, S.V. Side effects of BNT162b2 mRNA COVID-19 vaccine: A randomized, cross-sectional study with detailed self-reported symptoms from healthcare workers. *Int. J. Infect. Dis.* **2021**, *106*, 376–381. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Side+effects+of+BNT162b2+mRNA+COVID-19+vaccine:+A+randomized,+cross-sectional+study+with+detailed+self-reported+symptoms+from+healthcare+workers&author=Kadali,+R.A.K.&author=Janagama,+R.&author=Peruru,+S.&author=Malayala,+S.V.&publication_year=2021&journal=Int.+J.+Infect.+Dis.&volume=106&pages=376%E2%80%93381&doi=10.1016/j.ijid.2021.04.047&pmid=33866000)] [CrossRef] (<https://doi.org/10.1016/j.ijid.2021.04.047>) [PubMed] (<https://www.ncbi.nlm.nih.gov/pubmed/33866000>)]
14. Parkash, O.; Sharko, A.; Farooqi, A.; Ying, G.W.; Sura, P. Acute Pancreatitis: A Possible Side Effect of COVID-19 Vaccine. *Cureus* **2021**, *13*, e14741. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Acute+Pancreatitis:+A+Possible+Side+Effect+of+COVID-19+Vaccine&author=Parkash,+O.&author=Sharko,+A.&author=Farooqi,+A.&author=Ying,+G.W.&author=Sura,+P.&publication_year=2021&journal=Cureus&volume=13&pages=e14741&doi=10.7759/cureus.14741&pmid=34084669)] [CrossRef] (<https://doi.org/10.7759/cureus.14741>) [PubMed] (<https://www.ncbi.nlm.nih.gov/pubmed/34084669>)]

15. Mazzatenta, C.; Piccolo, V.; Pace, G.; Romano, I.; Argenziano, G.; Bassi, A. Purpuric lesions on the eyelids developed after BNT162b2 mRNA COVID-19 vaccine: Another piece of SARS-CoV-2 skin puzzle? *J. Eur. Acad. Dermatol. Venereol.* **2021**, *35*, e543–e545. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Purpuric+lesions+on+the+eyelids+developed+after+BNT162b2+mRNA+COVID-19+vaccine:+Another+piece+of+SARS-CoV-2+skin+puzzle?&author=Mazzatenta,+C.&author=Piccolo,+V.&author=Pace,+G.&author=Romano,+I.&author=Argenziano,+G.&author=Bassi,+A.&publication_year=2021&journal=J.+Eur.+Acad.+Dermatol.+Venereol.&volume=35&pages=e543%E2%80%93e545&doi=10.1111/jdv.17340)] [CrossRef (<https://doi.org/10.1111/jdv.17340>)]
16. Lee, E.J.; Cines, D.B.; Gernsheimer, T.; Kessler, C.; Michel, M.; Tarantino, M.D.; Semple, J.W.; Arnold, D.M.; Godeau, B.; Lambert, M.P.; et al. Thrombocytopenia following Pfizer and Moderna SARS-CoV-2 vaccination. *Am. J. Hematol.* **2021**, *96*, 534–537. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Thrombocytopenia+following+Pfizer+and+Moderna+SARS-CoV-2+vaccination&author=Lee,+E.J.&author=Cines,+D.B.&author=Gernsheimer,+T.&author=Kessler,+C.&author=Michel,+M.&author=Tarantino,+M.D.&author=Semple,+J.W.&author=Arnold,+D.M.&author=Godeau,+B.&author=Lambert,+M.P.&publication_year=2021&journal=Am.+J.+Hematol.&volume=96&pages=534%E2%80%93537&doi=10.1002/ajh.26132)] [CrossRef (<https://doi.org/10.1002/ajh.26132>)]
17. Ishay, Y.; Kenig, A.; Tsemach-Toren, T.; Amer, R.; Rubin, L.; HersHKovitz, Y.; Kharouf, F. Autoimmune phenomena following SARS-CoV-2 vaccination. *Int. Immunopharmacol.* **2021**, *99*, 107970. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Autoimmune+phenomena+following+SARS-CoV-2+vaccination&author=Ishay,+Y.&author=Kenig,+A.&author=Tsemach-Toren,+T.&author=Amer,+R.&author=Rubin,+L.&author=HersHKovitz,+Y.&author=Kharouf,+F.&publication_year=2021&journal=Int.+Immunopharmacol.&volume=99&pages=107970&doi=10.1016/j.intimp.2021.107970)] [CrossRef (<https://doi.org/10.1016/j.intimp.2021.107970>)]

18. Das, B.B.; Kohli, U.; Ramachandran, P.; Nguyen, H.H.; Greil, G.; Hussain, T.; Tandon, A.; Kane, C.; Avula, S.; Duru, C.; et al. Myopericarditis following mRNA COVID-19 Vaccination in Adolescents 12 through 18 Years of Age. *J. Pediatr.* **2021**, *238*, 26–32.e1. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Myopericarditis+following+mRNA+COVID-19+Vaccination+in+Adolescents+12+through+18+Years+of+Age&author=Das,+B.B.+author=Kohli,+U.+author=Ramachandran,+P.+author=Nguyen,+H.H.+author=Greil,+G.+author=Hussain,+T.+author=Tandon,+A.+author=Kane,+C.+author=Avula,+S.+author=Duru,+C.+publication_year=2021&journal=J.+Pediatr.&volume=238&pages=26%E2%80%9332.e1&doi=10.1016/j.jpeds.2021.07.044)] [CrossRef (<https://doi.org/10.1016/j.jpeds.2021.07.044>)]
19. McLaurin-Jiang, S.; Garner, C.D.; Krutsch, K.; Hale, T.W. Maternal and Child Symptoms Following COVID-19 Vaccination Among Breastfeeding Mothers. *Breastfeed. Med.* **2021**, *16*, 702–709. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Maternal+and+Child+Symptoms+Following+COVID-19+Vaccination+Among+Breastfeeding+Mothers&author=McLaurin-Jiang,+S.+author=Garner,+C.D.+author=Krutsch,+K.+author=Hale,+T.W.+publication_year=2021&journal=Breastfeed.+Med.&volume=16&pages=702%E2%80%93709&doi=10.1089/bfm.2021.0079)] [CrossRef (<https://doi.org/10.1089/bfm.2021.0079>)]
20. Barda, N.; Dagan, N.; Ben-Shlomo, Y.; Kepten, E.; Waxman, J.; Ohana, R.; Hernan, M.A.; Lipsitch, M.; Kohane, I.; Netzer, D.; et al. Safety of the BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Setting. *N. Engl. J. Med.* **2021**, *385*, 1078–1090. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Safety+of+the+BNT162b2+mRNA+Covid-19+Vaccine+in+a+Nationwide+Setting&author=Barda,+N.+author=Dagan,+N.+author=Ben-Shlomo,+Y.+author=Kepten,+E.+author=Waxman,+J.+author=Ohana,+R.+author=Hernan,+M.A.+author=Lipsitch,+M.+author=Kohane,+I.+author=Netzer,+D.+publication_year=2021&journal=N.+Engl.+J.+Med.&volume=385&pages=1078%E2%80%931090&doi=10.1056/NEJMoa2110475)] [CrossRef (<https://doi.org/10.1056/NEJMoa2110475>)]

21. Baden, L.R.; El Sahly, H.M.; Essink, B.; Kotloff, K.; Frey, S.; Novak, R.; Diemert, D.; Spector, S.A.; Rouphael, N.; Creech, C.B.; et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N. Engl. J. Med.* **2021**, *384*, 403–416. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Efficacy+and+Safety+of+the+mRNA-1273+SARS-CoV-2+Vaccine&author=Baden,+L.R.&author=El+Sahly,+H.M.&author=Essink,+B.&author=Kotloff,+K.&author=Frey,+S.&author=Novak,+R.&author=Diemert,+D.&author=Spector,+S.A.&author=Rouphael,+N.&author=Creech,+C.B.&publication_year=2021&journal=N.+Engl.+J.+Med.&volume=384&pages=403%E2%80%93416&doi=10.1056/NEJMoa2035389)] [CrossRef (<https://doi.org/10.1056/NEJMoa2035389>)]
22. Sadoff, J.; Gray, G.; Vandebosch, A.; Cardenas, V.; Shukarev, G.; Grinsztejn, B.; Goepfert, P.A.; Truyers, C.; Fennema, H.; Spiessens, B.; et al. Safety and Efficacy of Single-Dose Ad26.COV2.S Vaccine against Covid-19. *N. Engl. J. Med.* **2021**, *384*, 2187–2201. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Safety+and+Efficacy+of+Single-Dose+Ad26.COV2.S+Vaccine+against+Covid-19&author=Sadoff,+J.&author=Gray,+G.&author=Vandebosch,+A.&author=Cardenas,+V.&author=Shukarev,+G.&author=Grinsztejn,+B.&author=Goepfert,+P.A.&author=Truyers,+C.&author=Fennema,+H.&author=Spiessens,+B.&publication_year=2021&journal=N.+Engl.+J.+Med.&volume=384&pages=2187%E2%80%932201&doi=10.1056/NEJMoa2101544&pmid=33882225)] [CrossRef (<https://doi.org/10.1056/NEJMoa2101544>)] [PubMed (<https://www.ncbi.nlm.nih.gov/pubmed/33882225>)]
23. Eichinger, S.; Warkentin, T.E.; Greinacher, A. Thrombotic Thrombocytopenia after ChAdOx1 nCoV-19 Vaccination. Reply. *N. Engl. J. Med.* **2021**, *385*, e11. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Thrombotic+Thrombocytopenia+after+ChAdOx1+nCoV-19+Vaccination.+Reply&author=Eichinger,+S.&author=Warkentin,+T.E.&author=Greinacher,+A.&publication_year=2021&journal=N.+Engl.+J.+Med.&volume=385&pages=e11&doi=10.1056/NEJMc2107227&pmid=34133853)] [CrossRef (<https://doi.org/10.1056/NEJMc2107227>)] [PubMed (<https://www.ncbi.nlm.nih.gov/pubmed/34133853>)]

24. Doroftei, B.; Ciobica, A.; Ilie, O.D.; Maftai, R.; Ilea, C. Mini-Review Discussing the Reliability and Efficiency of COVID-19 Vaccines. *Diagnostics* **2021**, *11*, 579. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Mini-Review+Discussing+the+Reliability+and+Efficiency+of+COVID-19+Vaccines&author=Doroftei,+B.&author=Ciobica,+A.&author=Ilie,+O.D.&author=Maftai,+R.&author=Ilea,+C.&publication_year=2021&journal=Diagnostics&volume=11&pages=579&doi=10.3390/diagnostics11040579)] [CrossRef (https://doi.org/10.3390/diagnostics11040579)]

25. Zhang, L.; Richards, A.; Barrasa, M.I.; Hughes, S.H.; Young, R.A.; Jaenisch, R. Reverse-transcribed SARS-CoV-2 RNA can integrate into the genome of cultured human cells and can be expressed in patient-derived tissues. *Proc. Natl. Acad. Sci. USA* **2021**, *118*, e2105968118. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Reverse-transcribed+SARS-CoV-2+RNA+can+integrate+into+the+genome+of+cultured+human+cells+and+can+be+expressed+in+patient-derived+tissues&author=Zhang,+L.&author=Richards,+A.&author=Barrasa,+M.I.&author=Hughes,+S.H.&author=Young,+R.A.&author=Jaenisch,+R.&publication_year=2021&journal=Proc.+Natl.+Acad.+Sci.+USA&volume=118&pages=e2105968118&doi=10.1073/pnas.2105968118&pmid=33958444)] [CrossRef (https://doi.org/10.1073/pnas.2105968118)] [PubMed (https://www.ncbi.nlm.nih.gov/pubmed/33958444)]

26. Available online: https://www.ema.europa.eu/en/documents/assessment-report/comirnaty-epar-public-assessment-report_en.pdf (https://www.ema.europa.eu/en/documents/assessment-report/comirnaty-epar-public-assessment-report_en.pdf) (accessed on 24 February 2022).

27. Tanaka, H.; Takata, N.; Sakurai, Y.; Yoshida, T.; Inoue, T.; Tamagawa, S.; Nakai, Y.; Tange, K.; Yoshioka, H.; Maeki, M.; et al. Delivery of Oligonucleotides Using a Self-Degradable Lipid-Like Material. *Pharmaceutics* **2021**, *13*, 544. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Delivery+of+Oligonucleotides+Using+a+Self-Degradable+Lipid-Like+Material&author=Tanaka,+H.&author=Takata,+N.&author=Sakurai,+Y.&author=Yoshida,+T.&author=Inoue,+T.&author=Tamagawa,+S.&author=Nakai,+Y.&author=Tange,+K.&author=Yoshioka,+H.&author=Maeki,+M.&publication_year=2021&journal=Pharmaceutics&volume=13&pages=544&doi=10.3390/pharmaceutics13040544)] [CrossRef (https://doi.org/10.3390/pharmaceutics13040544)]

We use cookies to enhance your browsing experience. By clicking OK, you agree to our use of cookies. We also use cookies for analytics, to improve our website, and for marketing purposes. You can manage your cookie preferences at any time. Read more about our cookies (https://doi.org/10.3390/pharmaceutics13040544)]

28. Sedic, M.; Senn, J.J.; Lynn, A.; Laska, M.; Smith, M.; Platz, S.J.; Bolen, J.; Hoge, S.; Bulychhev, A.; Jacquinet, E.; et al. Safety Evaluation of Lipid Nanoparticle-Formulated Modified mRNA in the Sprague-Dawley Rat and Cynomolgus Monkey. *Vet. Pathol.* **2018**, *55*, 341–354. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Safety+Evaluation+of+Lipid+Nanoparticle-Formulated+Modified+mRNA+in+the+Sprague-Dawley+Rat+and+Cynomolgus+Monkey&author=Sedic,+M.&author=Senn,+J.J.&author=Lynn,+A.&author=Laska,+M.&author=Smith,+M.&author=Platz,+S.J.&author=Bolen,+J.&author=Hoge,+S.&author=Bulychhev,+A.&author=Jacquinet,+E.&publication_year=2018&journal=Vet.+Pathol.&volume=55&pages=341%E2%80%93354&doi=10.1177/0300985817738095)] [CrossRef (<https://doi.org/10.1177/0300985817738095>)]

29. Sato, Y.; Matsui, H.; Yamamoto, N.; Sato, R.; Munakata, T.; Kohara, M.; Harashima, H. Highly specific delivery of siRNA to hepatocytes circumvents endothelial cell-mediated lipid nanoparticle-associated toxicity leading to the safe and efficacious decrease in the hepatitis B virus. *J. Control. Release* **2017**, *266*, 216–225. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Highly+specific+delivery+of+siRNA+to+hepatocytes+circumvents+endothelial+cell-mediated+lipid+nanoparticle-associated+toxicity+leading+to+the+safe+and+efficacious+decrease+in+the+hepatitis+B+virus&author=Sato,+Y.&author=Matsui,+H.&author=Yamamoto,+N.&author=Sato,+R.&author=Munakata,+T.&author=Kohara,+M.&author=Harashima,+H.&publication_year=2017&journal=J.+Control.+Release&volume=266&pages=216%E2%80%93225&doi=10.1016/j.jconrel.2017.09.044)] [CrossRef (<https://doi.org/10.1016/j.jconrel.2017.09.044>)]

30. Heidel, J.D.; Yu, Z.; Liu, J.Y.; Rele, S.M.; Liang, Y.; Zeidan, R.K.; Kornbrust, D.J.; Davis, M.E. Administration in non-human primates of escalating intravenous doses of targeted nanoparticles containing ribonucleotide reductase subunit M2 siRNA. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 5715–5721. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Administration+in+non-human+primates+of+escalating+intravenous+doses+of+targeted+nanoparticles+containing+ribonucleotide+reductase+subunit+M2+siRNA&author=Heidel,+J.D.&author=Yu,+Z.&author=Liu,+J.Y.&author=Rele,+S.M.&author=Liang,+Y.&author=Zeidan,+R.K.&author=Kornbrust,+D.J.&author=Davis,+M.E.&publication_year=2007&journal=Proc.+Natl.+Acad.+Sci.+USA&volume=104&pages=5715%E2%80%935721&doi=10.1073/pnas.0701458104)] [CrossRef (<https://doi.org/10.1073/pnas.0701458104>)]

We use cookies on our website to ensure you get the best experience.

Read more about our cookies here ([about privacy](#)).

MDPI

31 Available online: <https://www.cvdvaccine-us.com/> (<https://www.cvdvaccine-us.com/>) (accessed on 24 February 2022).

32. Available online: [http://bridgeslab.sph.umich.edu/protocols/index.php/Preparation_of_Tail_Samples_\(for_Genotyping\)](http://bridgeslab.sph.umich.edu/protocols/index.php/Preparation_of_Tail_Samples_(for_Genotyping)) ([http://bridgeslab.sph.umich.edu/protocols/index.php/Preparation_of_Tail_Samples_\(for_Genotyping\)](http://bridgeslab.sph.umich.edu/protocols/index.php/Preparation_of_Tail_Samples_(for_Genotyping))) (accessed on 24 February 2022).
33. Gallud, A.; Munson, M.J.; Liu, K.; Idstrom, A.; Barriga, H.M.; Tabaei, S.; Aliakbarinodehi, N.; Ojansivu, M.; Lubart, Q.; Douth, J.J.; et al. Time evolution of PEG-shedding and serum protein coronation determines the cell uptake kinetics and delivery of lipid nanoparticle. *bioRxiv* **2021**. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Time+evolution+of+PEG-shedding+and+serum+protein+coronation+determines+the+cell+uptake+kinetics+and+delivery+of+lipid+nanoparticle&author=Gallud,+A.&author=Munson,+M.J.&author=Liu,+K.&author=Idstrom,+A.&author=Barriga,+H.M.&author=Tabaei,+S.&author=Aliakbarinodehi,+N.&author=Ojansivu,+M.&author=Lubart,+Q.&author=Douth,+J.J.&publication_year=2021&journal=bioRxiv&doi=10.1101/2021.08.20.457104)] [CrossRef (<https://doi.org/10.1101/2021.08.20.457104>)]
34. World Health Organization Messenger RNA Encoding the Full-Length SARS-CoV-2 Spike Glycoprotein. 2020. Available online: <https://web.archive.org/web/20210105162941/https://mednet-communities.net/inn/db/media/docs/11889.doc> (<https://web.archive.org/web/20210105162941/https://mednet-communities.net/inn/db/media/docs/11889.doc>) (accessed on 24 February 2022).
35. Mita, P.; Wudzinska, A.; Sun, X.; Andrade, J.; Nayak, S.; Kahler, D.J.; Badri, S.; LaCava, J.; Ueberheide, B.; Yun, C.Y.; et al. LINE-1 protein localization and functional dynamics during the cell cycle. *Elife* **2018**, *7*, e30058. [Google Scholar (https://scholar.google.com/scholar_lookup?title=LINE-1+protein+localization+and+functional+dynamics+during+the+cell+cycle&author=Mita,+P.&author=Wudzinska,+A.&author=Sun,+X.&author=Andrade,+J.&author=Nayak,+S.&author=Kahler,+D.J.&author=Badri,+S.&author=LaCava,+J.&author=Ueberheide,+B.&author=Yun,+C.Y.&publication_year=2018&journal=Elife&volume=7&pages=e30058&doi=10.7554/eLife.30058)] [CrossRef (<https://doi.org/10.7554/eLife.30058>)]

We use cookies on our website to ensure you get the best experience.
Read more about our cookies [here \(/about/privacy\)](#).

[Accept \(/accept_cookies\)](#)

36 Sato, Y.; Kinami, Y.; Hashiba, K.; Harashima, H. Different kinetics for the hepatic uptake of lipid nanoparticles between the apolipoprotein E/low density lipoprotein receptor and the N-acetyl-d-galactosamine/asialoglycoprotein receptor pathway. *J. Control. Release* **2020**, *322*, 217–226. [Google Scholar

(https://scholar.google.com/scholar_lookup?title=Different+kinetics+for+the+hepatic+uptake+of+lipid+nanoparticles+between+the+apolipoprotein+E/low+density+lipoprotein+receptor+and+the+N-acetyl-d-galactosamine/asialoglycoprotein+receptor+pathway&author=Sato,+Y.&author=Kinami,+Y.&author=Hashiba,+K.&author=Harashima,+H.&publication_year=2020&journal=J.+Control.+Release&volume=322&pages=217%E2%80%93226&doi=10.1016/j.jconrel.2020.03.006)] [CrossRef

(<https://doi.org/10.1016/j.jconrel.2020.03.006>)]

37. Vogel, A.B.; Kanevsky, I.; Che, Y.; Swanson, K.A.; Muik, A.; Vormehr, M.; Kranz, L.M.; Walzer, K.C.; Hein, S.; Guler, A.; et al. BNT162b vaccines protect rhesus macaques from SARS-CoV-2. *Nature* **2021**, *592*, 283–289. [Google Scholar

(https://scholar.google.com/scholar_lookup?title=BNT162b+vaccines+protect+rhesus+macaques+from+SARS-CoV-2&author=Vogel,+A.B.&author=Kanevsky,+I.&author=Che,+Y.&author=Swanson,+K.A.&author=Muik,+A.&author=Vormehr,+M.&author=Kranz,+L.M.&author=Walzer,+K.C.&author=Hein,+S.&author=Guler,+A.&publication_year=2021&journal=Nature&volume=592&pages=283%E2%80%93289&doi=10.1038/s41586-021-03275-y&pmid=33524990)] [CrossRef (<https://doi.org/10.1038/s41586-021-03275-y>)] [PubMed (<https://www.ncbi.nlm.nih.gov/pubmed/33524990>)]

38. Bahl, K.; Senn, J.J.; Yuzhakov, O.; Bulychhev, A.; Brito, L.A.; Hassett, K.J.; Laska, M.E.; Smith, M.; Almarsson, O.; Thompson, J.; et al. Preclinical and Clinical Demonstration of Immunogenicity by mRNA Vaccines against H10N8 and H7N9 Influenza Viruses. *Mol. Ther.* **2017**, *25*, 1316–1327. [Google Scholar

(https://scholar.google.com/scholar_lookup?title=Preclinical+and+Clinical+Demonstration+of+Immunogenicity+by+mRNA+Vaccines+against+H10N8+and+H7N9+Influenza+Viruses&author=Bahl,+K.&author=Senn,+J.J.&author=Yuzhakov,+O.&author=Bulychhev,+A.&author=Brito,+L.A.&author=Hassett,+K.J.&author=Laska,+M.E.&author=Smith,+M.&author=Almarsson,+O.&author=Thompson,+J.&publication_year=2017&journal=Mol.+Ther.&volume=25&pages=1316%E2%80%931327&doi=10.1016/j.ymthe.2017.03.035&pmid=28457665)] [CrossRef (<https://doi.org/10.1016/j.ymthe.2017.03.035>)] [PubMed

We use cookies on our website to ensure you get the best experience.

Read more about our cookies [here](#) (about privacy).

39. Bril, F.; Al Diffalha, S.; Dean, M.; Fettig, D.M. Autoimmune hepatitis developing after coronavirus disease 2019 (COVID-19) vaccine: Causality or casualty? *J. Hepatol.* **2021**, *75*, 222–224. [Google Scholar ([https://scholar.google.com/scholar_lookup?title=Autoimmune+hepatitis+developing+after+coronavirus+disease+2019+\(COVID-19\)+vaccine:+Causality+or+casualty?&author=Bril,+F.&author=Al+Diffalha,+S.&author=Dean,+M.&author=Fettig,+D.M.&publication_year=2021&journal=J.+Hepatol.&volume=75&pages=222%E2%80%93224&doi=10.1016/j.jhep.2021.04.003](https://scholar.google.com/scholar_lookup?title=Autoimmune+hepatitis+developing+after+coronavirus+disease+2019+(COVID-19)+vaccine:+Causality+or+casualty?&author=Bril,+F.&author=Al+Diffalha,+S.&author=Dean,+M.&author=Fettig,+D.M.&publication_year=2021&journal=J.+Hepatol.&volume=75&pages=222%E2%80%93224&doi=10.1016/j.jhep.2021.04.003))] [CrossRef (<https://doi.org/10.1016/j.jhep.2021.04.003>)]

40. Kazazian, H.H., Jr.; Moran, J.V. Mobile DNA in Health and Disease. *N. Engl. J. Med.* **2017**, *377*, 361–370. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Mobile+DNA+in+Health+and+Disease&author=Kazazian,+H.H.,+Jr.&author=Moran,+J.V.&publication_year=2017&journal=N.+Engl.+J.+Med.&volume=377&pages=361%E2%80%93370&doi=10.1056/NEJMra1510092&pmid=28745987)] [CrossRef (<https://doi.org/10.1056/NEJMra1510092>)] [PubMed (<https://www.ncbi.nlm.nih.gov/pubmed/28745987>)]

41. Coffin, J.M.; Fan, H. The Discovery of Reverse Transcriptase. *Annu. Rev. Virol.* **2016**, *3*, 29–51. [Google Scholar (https://scholar.google.com/scholar_lookup?title=The+Discovery+of+Reverse+Transcriptase&author=Coffin,+J.M.&author=Fan,+H.&publication_year=2016&journal=Annu.+Rev.+Virol.&volume=3&pages=29%E2%80%9351&doi=10.1146/annurev-virology-110615-035556)] [CrossRef (<https://doi.org/10.1146/annurev-virology-110615-035556>)]

42. Lander, E.S.; Linton, L.M.; Birren, B.; Nusbaum, C.; Zody, M.C.; Baldwin, J.; Devon, K.; Dewar, K.; Doyle, M.; FitzHugh, W.; et al. Initial sequencing and analysis of the human genome. *Nature* **2001**, *409*, 860–921. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Initial+sequencing+and+analysis+of+the+human+genome&author=Lander,+E.S.&author=Linton,+L.M.&author=Birren,+B.&author=Nusbaum,+C.&author=Zody,+M.C.&author=Baldwin,+J.&author=Devon,+K.&author=Dewar,+K.&author=Doyle,+M.&author=FitzHugh,+W.&publication_year=2001&journal=Nature&volume=409&pages=860%E2%80%93921&doi=10.1038/35057062)] [CrossRef (<https://doi.org/10.1038/35057062>)]

We use cookies on our website to ensure you get the best experience.
Read more about our cookies [here \(/about/privacy\)](#).

[Accept \(/accept_cookies\)](#)

43. Ostertag, E.M.; Goodier, J.L.; Zhang, Y.; Kazazian, H.H., Jr. SVA elements are nonautonomous retrotransposons that cause disease in humans. *Am. J. Hum. Genet.* **2003**, *73*, 1444–1451. [Google Scholar

(https://scholar.google.com/scholar_lookup?

[title=SVA+elements+are+nonautonomous+retrotransposons+that+cause+disease+in+humans&author=Ostertag,+E.M.&author=Goodier,+J.L.&author=Zhang,+Y.&author=Kazazian,+H.H.,+Jr.&publication_year=2003&journal=Am.+J.+Hum.+Genet.&volume=73&pages=1444%E2%80%931451&doi=10.1086/380207](https://scholar.google.com/scholar_lookup?title=SVA+elements+are+nonautonomous+retrotransposons+that+cause+disease+in+humans&author=Ostertag,+E.M.&author=Goodier,+J.L.&author=Zhang,+Y.&author=Kazazian,+H.H.,+Jr.&publication_year=2003&journal=Am.+J.+Hum.+Genet.&volume=73&pages=1444%E2%80%931451&doi=10.1086/380207)] [CrossRef (<https://doi.org/10.1086/380207>)]

44. Hancks, D.C.; Kazazian, H.H., Jr. Active human retrotransposons: Variation and disease. *Curr. Opin. Genet. Dev.* **2012**, *22*, 191–203. [Google Scholar (https://scholar.google.com/scholar_lookup?

[title=Active+human+retrotransposons:+Variation+and+disease&author=Hancks,+D.C.&author=Kazazian,+H.H.,+Jr.&publication_year=2012&journal=Curr.+Opin.+Genet.+Dev.&volume=22&pages=191%E2%80%93203&doi=10.1016/j.gde.2012.02.006](https://scholar.google.com/scholar_lookup?title=Active+human+retrotransposons:+Variation+and+disease&author=Hancks,+D.C.&author=Kazazian,+H.H.,+Jr.&publication_year=2012&journal=Curr.+Opin.+Genet.+Dev.&volume=22&pages=191%E2%80%93203&doi=10.1016/j.gde.2012.02.006)] [CrossRef (<https://doi.org/10.1016/j.gde.2012.02.006>)]

45. Jones, R.B.; Song, H.; Xu, Y.; Garrison, K.E.; Buzdin, A.A.; Anwar, N.; Hunter, D.V.; Mujib, S.; Mihajlovic, V.; Martin, E.; et al. LINE-1 retrotransposable element DNA accumulates in HIV-1-infected cells. *J. Virol.* **2013**, *87*, 13307–13320. [Google Scholar (https://scholar.google.com/scholar_lookup?title=LINE-1+retrotransposable+element+DNA+accumulates+in+HIV-1-infected+cells&author=Jones,+R.B.&author=Song,+H.&author=Xu,+Y.&author=Garrison,+K.E.&author=Buzdin,+A.A.&author=Anwar,+N.&author=Hunter,+D.V.&author=Mujib,+S.&author=Mihajlovic,+V.&author=Martin,+E.&publication_year=2013&journal=J.+Virol.&volume=87&pages=13307%E2%80%9313320&doi=10.1128/JVI.02257-13)] [CrossRef (<https://doi.org/10.1128/JVI.02257-13>)]

46. Macchietto, M.G.; Langlois, R.A.; Shen, S.S. Virus-induced transposable element expression up-regulation in human and mouse host cells. *Life Sci. Alliance* **2020**, *3*, e201900536. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Virus-induced+transposable+element+expression+up-regulation+in+human+and+mouse+host+cells&author=Macchietto,+M.G.&author=Langlois,+R.A.&author=Shen,+S.S.&publication_year=2020&journal=Life+Sci.+Alliance&volume=3&pages=e201900536&doi=10.26508/lsa.201900536&pmid=31964680)] [CrossRef (<https://doi.org/10.26508/lsa.201900536>)] [PubMed (<https://pubmed.ncbi.nlm.nih.gov/31964680/>)]

We use cookies to enhance your browsing experience.

Read more about our cookies [here](#) ([/about/privacy](#)).

47. Yin, Y.; Liu, X.Z.; He, X.; Zhou, L.Q. Exogenous Coronavirus Interacts With Endogenous Retrotransposon in Human Cells. *Front. Cell Infect. Microbiol.* **2021**, *11*, 609160. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Exogenous+Coronavirus+Interacts+With+Endogenous+Retrotransposon+in+Human+Cells&author=Yin,+Y.&author=Liu,+X.Z.&author=He,+X.&author=Zhou,+L.Q.&publication_year=2021&journal=Front.+Cell+Infect.+Microbiol.&volume=11&pages=609160&doi=10.3389/fcimb.2021.609160&pmid=33732659)] [CrossRef (<https://doi.org/10.3389/fcimb.2021.609160>)] [PubMed (<https://www.ncbi.nlm.nih.gov/pubmed/33732659>)]

48. Belancio, V.P.; Roy-Engel, A.M.; Deininger, P. The impact of multiple splice sites in human L1 elements. *Gene* **2008**, *411*, 38–45. [Google Scholar (https://scholar.google.com/scholar_lookup?title=The+impact+of+multiple+splice+sites+in+human+L1+elements&author=Belancio,+V.P.&author=Roy-Engel,+A.M.&author=Deininger,+P.&publication_year=2008&journal=Gene&volume=411&pages=38%E2%80%9345&doi=10.1016/j.gene.2007.12.022&pmid=18261861)] [CrossRef (<https://doi.org/10.1016/j.gene.2007.12.022>)] [PubMed (<https://www.ncbi.nlm.nih.gov/pubmed/18261861>)]

49. Dai, L.; Taylor, M.S.; O'Donnell, K.A.; Boeke, J.D. Poly(A) binding protein C1 is essential for efficient L1 retrotransposition and affects L1 RNP formation. *Mol. Cell Biol.* **2012**, *32*, 4323–4336. [Google Scholar ([https://scholar.google.com/scholar_lookup?title=Poly\(A\)+binding+protein+C1+is+essential+for+efficient+L1+retrotransposition+and+affects+L1+RNP+formation&author=Dai,+L.&author=Taylor,+M.S.&author=O'Donnell,+K.A.&author=Boeke,+J.D.&publication_year=2012&journal=Mol.+Cell+Biol.&volume=32&pages=4323%E2%80%934336&doi=10.1128/MCB.06785-11](https://scholar.google.com/scholar_lookup?title=Poly(A)+binding+protein+C1+is+essential+for+efficient+L1+retrotransposition+and+affects+L1+RNP+formation&author=Dai,+L.&author=Taylor,+M.S.&author=O'Donnell,+K.A.&author=Boeke,+J.D.&publication_year=2012&journal=Mol.+Cell+Biol.&volume=32&pages=4323%E2%80%934336&doi=10.1128/MCB.06785-11))] [CrossRef (<https://doi.org/10.1128/MCB.06785-11>)]

50. Servant, G.; Strevi, V.A.; Derbes, R.S.; Wijetunge, M.I.; Neeland, M.; White, T.B.; Belancio, V.P.; Roy-Engel, A.M.; Deininger, P.L. The Nucleotide Excision Repair Pathway Limits L1 Retrotransposition. *Genetics* **2017**, *205*, 139–153. [Google Scholar (https://scholar.google.com/scholar_lookup?title=The+Nucleotide+Excision+Repair+Pathway+Limits+L1+Retrotransposition&author=Servant,+G.&author=Strevi,+V.A.&author=Derbes,+R.S.&author=Wijetunge,+M.I.&author=Neeland,+M.&author=White,+T.B.&author=Belancio,+V.P.&author=Roy-Engel,+A.M.&author=Deininger,+P.L.&publication_year=2017&journal=Genetics&volume=205&pages=139%E2%80%93153&doi=10.1534/genetics.116.188680)] [CrossRef (<https://doi.org/10.1534/genetics.116.188680>)]

We use cookies on our website to ensure you get the best experience.

Read more about our cookies here ([https://www.mdpi.com/privacy-policy/accept_cookies](#))

51. Guo, H.; Chitiprolu, M.; Gagnon, D.; Meng, L.; Perez-Iratxeta, C.; Lagace, D.; Gibbings, D. Autophagy supports genomic stability by degrading retrotransposon RNA. *Nat. Commun.* **2014**, *5*, 5276. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Autophagy+supports+genomic+stability+by+degrading+retrotransposon+RNA&author=Guo,+H.&author=Chitiprolu,+M.&author=Gagnon,+D.&author=Meng,+L.&author=Perez-Iratxeta,+C.&author=Lagace,+D.&author=Gibbings,+D.&publication_year=2014&journal=Nat.+Commun.&volume=5&pages=5276&doi=10.1038/ncomms6276)] [CrossRef (https://doi.org/10.1038/ncomms6276)]
52. Xie, Y.; Mates, L.; Ivics, Z.; Izsvak, Z.; Martin, S.L.; An, W. Cell division promotes efficient retrotransposition in a stable L1 reporter cell line. *Mob. DNA* **2013**, *4*, 10. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Cell+division+promotes+efficient+retrotransposition+in+a+stable+L1+reporter+cell+line&author=Xie,+Y.&author=Mates,+L.&author=Ivics,+Z.&author=Izsvak,+Z.&author=Martin,+S.L.&author=An,+W.&publication_year=2013&journal=Mob.+DNA&volume=4&pages=10&doi=10.1186/1759-8753-4-10)] [CrossRef (https://doi.org/10.1186/1759-8753-4-10)]
53. Shi, X.; Seluanov, A.; Gorbunova, V. Cell divisions are required for L1 retrotransposition. *Mol. Cell Biol.* **2007**, *27*, 1264–1270. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Cell+divisions+are+required+for+L1+retrotransposition&author=Shi,+X.&author=Seluanov,+A.&author=Gorbunova,+V.&publication_year=2007&journal=Mol.+Cell+Biol.&volume=27&pages=1264%E2%80%931270&doi=10.1128/MCB.01888-06)] [CrossRef (https://doi.org/10.1128/MCB.01888-06)]
54. Goff, S.P. Host factors exploited by retroviruses. *Nat. Rev. Microbiol* **2007**, *5*, 253–263. [Google Scholar (https://scholar.google.com/scholar_lookup?title=Host+factors+exploited+by+retroviruses&author=Goff,+S.P.&publication_year=2007&journal=Nat.+Rev.+Microbiol&volume=5&pages=253%E2%80%93263&doi=10.1038/nrmicro1541)] [CrossRef (https://doi.org/10.1038/nrmicro1541)]
55. Suzuki, Y.; Craigie, R. The road to chromatin—Nuclear entry of retroviruses. *Nat. Rev. Microbiol.* **2007**, *5*, 187–196. [Google Scholar (https://scholar.google.com/scholar_lookup?title=The+road+to+chromatin%E2%80%93Nuclear+entry+of+retroviruses&author=Suzuki,+Y.&author=Craigie,+R.&publication_year=2007&journal=Nat.+Rev.+Microbiol.&volume=5&pages=187%E2%80%93196&doi=10.1038/nrmicro1579)] [CrossRef (https://doi.org/10.1038/nrmicro1579)]

56. Shi, J.; Wang, X.; Lyu, L.; Jiang, H.; Zhu, H.J. Comparison of protein expression between human livers and the hepatic cell lines HepG2, Hep3B, and Huh7 using SWATH and MRM-HR proteomics: Focusing on drug-metabolizing enzymes. *Drug Metab. Pharmacokinet.* **2018**, *33*, 133–140. [Google Scholar

([https://scholar.google.com/scholar_lookup?](https://scholar.google.com/scholar_lookup?title=Comparison+of+protein+expression+between+human+livers+and+the+hepatic+cell+lines+HepG2,+Hep3B,+and+Huh7+using+SWATH+and+MRM-HR+proteomics:+Focusing+on+drug-metabolizing+enzymes&author=Shi,+J.&author=Wang,+X.&author=Lyu,+L.&author=Jiang,+H.&author=Zhu,+H.J.&publication_year=2018&journal=Drug+Metab.+Pharmacokinet.&volume=33&pages=133%E2%80%93140&doi=10.1016/j.dmpk.2018.03.003&pmid=29610054)

[title=Comparison+of+protein+expression+between+human+livers+and+the+hepatic+cell+lines+HepG2,+Hep3B,+and+Huh7+using+SWATH+and+MRM-](https://scholar.google.com/scholar_lookup?title=Comparison+of+protein+expression+between+human+livers+and+the+hepatic+cell+lines+HepG2,+Hep3B,+and+Huh7+using+SWATH+and+MRM-HR+proteomics:+Focusing+on+drug-metabolizing+enzymes&author=Shi,+J.&author=Wang,+X.&author=Lyu,+L.&author=Jiang,+H.&author=Zhu,+H.J.&publication_year=2018&journal=Drug+Metab.+Pharmacokinet.&volume=33&pages=133%E2%80%93140&doi=10.1016/j.dmpk.2018.03.003&pmid=29610054)

[HR+proteomics:+Focusing+on+drug-metabolizing+enzymes&author=Shi,+J.&author=Wang,+X.&author=Lyu,+L.&author=Jiang,+H.&author=Zhu,+H.J.&publication_year=2018&journal=Drug+Metab.+Pharmacokinet.&volume=33&pages=133%E2%80%93140&doi=10.1016/j.dmpk.2018.03.003&pmid=29610054](https://scholar.google.com/scholar_lookup?title=Comparison+of+protein+expression+between+human+livers+and+the+hepatic+cell+lines+HepG2,+Hep3B,+and+Huh7+using+SWATH+and+MRM-HR+proteomics:+Focusing+on+drug-metabolizing+enzymes&author=Shi,+J.&author=Wang,+X.&author=Lyu,+L.&author=Jiang,+H.&author=Zhu,+H.J.&publication_year=2018&journal=Drug+Metab.+Pharmacokinet.&volume=33&pages=133%E2%80%93140&doi=10.1016/j.dmpk.2018.03.003&pmid=29610054)] [CrossRef

(<https://doi.org/10.1016/j.dmpk.2018.03.003>)]

[PubMed

(<https://www.ncbi.nlm.nih.gov/pubmed/29610054>)]

57. Kubo, S.; Seleme, M.C.; Soifer, H.S.; Perez, J.L.; Moran, J.V.; Kazazian, H.H., Jr.; Kasahara, N. L1 retrotransposition in nondividing and primary human somatic cells. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 8036–8041. [Google Scholar

([https://scholar.google.com/scholar_lookup?](https://scholar.google.com/scholar_lookup?title=L1+retrotransposition+in+nondividing+and+primary+human+somatic+cells&author=Kubo,+S.&author=Seleme,+M.C.&author=Soifer,+H.S.&author=Perez,+J.L.&author=Moran,+J.V.&author=Kazazian,+H.H.,+Jr.&author=Kasahara,+N.&publication_year=2006&journal=Proc.+Natl.+Acad.+Sci.+USA&volume=103&pages=8036%E2%80%938041&doi=10.1073/pnas.0601954103&pmid=16698926)

[title=L1+retrotransposition+in+nondividing+and+primary+human+somatic+cells&author=Kubo,+S.&author=Seleme,+M.C.&author=Soifer,+H.S.&author=Perez,+J.L.&author=Moran,+J.V.&author=Kazazian,+H.H.,+Jr.&author=Kasahara,+N.&publication_year=2006&journal=Proc.+Natl.+Acad.+Sci.+USA&volume=103&pages=8036%E2%80%938041&doi=10.1073/pnas.0601954103&pmid=16698926](https://scholar.google.com/scholar_lookup?title=L1+retrotransposition+in+nondividing+and+primary+human+somatic+cells&author=Kubo,+S.&author=Seleme,+M.C.&author=Soifer,+H.S.&author=Perez,+J.L.&author=Moran,+J.V.&author=Kazazian,+H.H.,+Jr.&author=Kasahara,+N.&publication_year=2006&journal=Proc.+Natl.+Acad.+Sci.+USA&volume=103&pages=8036%E2%80%938041&doi=10.1073/pnas.0601954103&pmid=16698926))]

[CrossRef (<https://doi.org/10.1073/pnas.0601954103>)]

[PubMed

(<https://www.ncbi.nlm.nih.gov/pubmed/16698926>)]

58. Macia, A.; Widmann, T.J.; Heras, S.R.; Ayllon, V.; Sanchez, L.; Benkaddour-Boumzaouad, M.; Munoz-Lopez, M.; Rubio, A.; Amador-Cubero, S.; Blanco-Jimenez, E.; et al. Engineered LINE-1 retrotransposition in nondividing human neurons. *Genome Res.* **2017**, *27*, 335–348. [Google Scholar

([https://scholar.google.com/scholar_lookup?title=Engineered+LINE-](https://scholar.google.com/scholar_lookup?title=Engineered+LINE-1+retrotransposition+in+nondividing+human+neurons&author=Macia,+A.&author=Widmann,+T.J.&author=Heras,+S.R.&author=Ayllon,+V.&author=Sanchez,+L.&author=Benkaddour-Boumzaouad,+M.&author=Munoz-Lopez,+M.&author=Rubio,+A.&author=Amador-Cubero,+S.&author=Blanco-Jimenez,+E.&publication_year=2017&journal=Genome+Res.&volume=27&pages=335%E2%80%93348&doi=10.1101/gr.206805.116&pmid=27965292)

[1+retrotransposition+in+nondividing+human+neurons&author=Macia,+A.&author=Widmann,+T.J.&author=Heras,+S.R.&author=Ayllon,+V.&author=Sanchez,+L.&author=Benkaddour-Boumzaouad,+M.&author=Munoz-](https://scholar.google.com/scholar_lookup?title=Engineered+LINE-1+retrotransposition+in+nondividing+human+neurons&author=Macia,+A.&author=Widmann,+T.J.&author=Heras,+S.R.&author=Ayllon,+V.&author=Sanchez,+L.&author=Benkaddour-Boumzaouad,+M.&author=Munoz-Lopez,+M.&author=Rubio,+A.&author=Amador-Cubero,+S.&author=Blanco-Jimenez,+E.&publication_year=2017&journal=Genome+Res.&volume=27&pages=335%E2%80%93348&doi=10.1101/gr.206805.116&pmid=27965292)

[Lopez,+M.&author=Rubio,+A.&author=Amador-Cubero,+S.&author=Blanco-](https://scholar.google.com/scholar_lookup?title=Engineered+LINE-1+retrotransposition+in+nondividing+human+neurons&author=Macia,+A.&author=Widmann,+T.J.&author=Heras,+S.R.&author=Ayllon,+V.&author=Sanchez,+L.&author=Benkaddour-Boumzaouad,+M.&author=Munoz-Lopez,+M.&author=Rubio,+A.&author=Amador-Cubero,+S.&author=Blanco-Jimenez,+E.&publication_year=2017&journal=Genome+Res.&volume=27&pages=335%E2%80%93348&doi=10.1101/gr.206805.116&pmid=27965292)

[Jimenez,+E.&publication_year=2017&journal=Genome+Res.&volume=27&pages=335%E2%80%93348&doi=10.1101/gr.206805.116&pmid=27965292](https://scholar.google.com/scholar_lookup?title=Engineered+LINE-1+retrotransposition+in+nondividing+human+neurons&author=Macia,+A.&author=Widmann,+T.J.&author=Heras,+S.R.&author=Ayllon,+V.&author=Sanchez,+L.&author=Benkaddour-Boumzaouad,+M.&author=Munoz-Lopez,+M.&author=Rubio,+A.&author=Amador-Cubero,+S.&author=Blanco-Jimenez,+E.&publication_year=2017&journal=Genome+Res.&volume=27&pages=335%E2%80%93348&doi=10.1101/gr.206805.116&pmid=27965292))] [CrossRef

(<https://doi.org/10.1101/gr.206805.116>)]

[PubMed

(<https://www.ncbi.nlm.nih.gov/pubmed/27965292>)]

[PubMed

(<https://www.ncbi.nlm.nih.gov/pubmed/27965292>)]

Accept (/accept_cookies)

We use cookies on our website to ensure you get the best experience.

Read more about our cookies here ([about/privacy](#)).

© 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>) (<https://creativecommons.org/licenses/by/4.0/>).

Curr. Issues Mol. Biol. (</journal/cimb>), EISSN 1467-3045, Published by MDPI [Disclaimer](#)
[RSS \(/rss/journal/cimb\)](/rss/journal/cimb) [Content Alert \(/journal/cimb/toc-alert\)](/journal/cimb/toc-alert)

← Further Information

[Article Processing Charges \(/apc\)](/apc)

[Pay an Invoice \(/about/payment\)](/about/payment)

[Open Access Policy \(/openaccess\)](/openaccess)

[Contact MDPI \(/about/contact\)](/about/contact)

[Jobs at MDPI \(https://careers.mdpi.com\)](https://careers.mdpi.com)

Guidelines

[For Authors \(/authors\)](/authors)

[For Reviewers \(/reviewers\)](/reviewers)

[For Editors \(/editors\)](/editors)

[For Librarians \(/librarians\)](/librarians)

[For Publishers \(/publishing_services\)](/publishing_services)

[For Societies \(/societies\)](/societies)

[For Conference Organizers \(/conference_organizers\)](/conference_organizers)

MDPI Initiatives

[Sciforum \(https://sciforum.net\)](https://sciforum.net)

[MDPI Books \(https://www.mdpi.com/books\)](https://www.mdpi.com/books)

[Preprints \(https://www.preprints.org\)](https://www.preprints.org)

[Scilit \(https://www.scilit.net\)](https://www.scilit.net)

[SciProfiles \(https://sciprofiles.com\)](https://sciprofiles.com)

[Encyclopedia \(https://encyclopedia.pub\)](https://encyclopedia.pub)

We use cookies on our website to ensure you get the best experience.

[JAMS \(https://jams.pub\)](https://jams.pub)

Read more about our cookies [here \(/about/privacy\)](/about/privacy).

[Proceedings Series \(/about/proceedings\)](/about/proceedings)

[Accept \(/accept_cookies\)](/accept_cookies)


Follow MDPI

LinkedIn (<https://www.linkedin.com/company/mdpi>)

MDPI ([/](#))

Facebook (<https://www.facebook.com/MDPIOpenAccessPublishing>)

Twitter (<https://twitter.com/MDPIOpenAccess>)



**Subscribe to receive issue release
notifications and newsletters from
MDPI journals**

Select options ▼

◀ Enter your email address... ▶

Subscribe

© 1996-2022 MDPI (Basel, Switzerland) unless otherwise stated

[Disclaimer](#)

[Terms and Conditions \(/about/terms-and-conditions\)](/about/terms-and-conditions)

[Privacy Policy \(/about/privacy\)](/about/privacy)

We use cookies on our website to ensure you get the best experience.

Read more about our cookies [here \(/about/privacy\)](/about/privacy).

[Accept \(/accept_cookies\)](/accept_cookies)